

MCL

Spark what's possible with Jaypirca® (pirtobrutinib)



Indication

Jaypirca is a prescription medicine used to treat adults with **mantle cell lymphoma (MCL)** that has come back or did not respond to previous treatment, and who have already received at least 2 treatments for their cancer, including a Bruton's tyrosine kinase (BTK) inhibitor medicine.

Jaypirca was approved for this indication based on how many people responded to treatment. Studies are ongoing to confirm the benefit of Jaypirca for this use.

It is not known if Jaypirca is safe and effective in children.

Select Important Safety Information

Warnings - Jaypirca may cause serious side effects, including:

Infections. Infections can happen during treatment with Jaypirca. These infections can be serious and may lead to death. Your doctor may prescribe vaccines and certain medicines if you have an increased risk of getting infections. Tell your doctor right away if you have fever, chills, weakness, flu-like symptoms, or any other signs of infection during treatment with Jaypirca.

Please see **Safety Summary** for Jaypirca on pages 8-10.

KEEP READING TO LEARN
MORE ABOUT JAYPIRCA

 **Jaypirca®**
pirtobrutinib 50,100 mg
tablets

A Lilly Medicine

What is Jaypirca?

Jaypirca is a prescription medicine used to treat adults with mantle cell lymphoma (MCL) that has come back or did not respond to previous treatment, and who have already received at least 2 treatments for their cancer, including a Bruton's tyrosine kinase (BTK) inhibitor medicine.

You may have taken BTK inhibitors before. BTK inhibitors stop, or inhibit, cancer cells from surviving and multiplying, which can slow the spread of certain cancers like MCL.

Jaypirca, a noncovalent BTK inhibitor, works differently than other BTK inhibitors you may have taken before.

For MCL, it was developed to work after you have taken at least 2 treatments, including a BTK inhibitor.



Not an actual patient.

Select Important Safety Information

Warnings - Jaypirca may cause serious side effects, including:

Bleeding problems (hemorrhage). Bleeding problems can happen during treatment with Jaypirca. They can be serious and may lead to death. Your risk of severe bleeding may increase if you are also taking a blood thinner medicine. Tell your doctor if you have any signs or symptoms of bleeding, including:

- blood in your stools or black stools (looks like tar)
- pink or brown urine
- unexpected bleeding, or bleeding that is severe or you cannot control
- vomit blood or vomit blood that looks like coffee grinds
- cough up blood or blood clots
- increased bruising
- dizziness
- weakness
- confusion
- changes in your speech
- headache that lasts a long time



Jaypirca[®]
pirtobrutinib 50, 100 mg
tablets


A Lilly Medicine

How to take Jaypirca

Take Jaypirca exactly as your healthcare provider tells you



Not an actual patient.



How do I take Jaypirca?

Take Jaypirca 1 time each day, at about the same time each day. Take Jaypirca with or without food. Swallow Jaypirca whole with water. Do not cut, crush, or chew the tablets.



What if I miss a dose?

If you miss a dose of Jaypirca, take it as soon as you remember on the same day. If it has been more than 12 hours, skip the missed dose and take your next dose on the next day at your usual time.



What if I experience side effects?

Your healthcare provider may tell you to decrease your dose, temporarily stop, or completely stop taking Jaypirca if you develop side effects. Do not change your dose or stop taking Jaypirca without talking to your healthcare provider first.

Select Important Safety Information

Warnings - Jaypirca may cause serious side effects, including:

Decrease in blood cell counts. Decrease in white blood cells, platelets, and red blood cells are common with Jaypirca, but can also be severe. This may increase your risk of infection, bleeding, and anemia. Your doctor should do blood tests regularly during treatment to monitor your counts.

Heart rhythm problems. Heart rhythm problems including atrial fibrillation and atrial flutter have happened in people treated with Jaypirca. Your risk for heart rhythm problems may be increased if you have high blood pressure or have had heart rhythm problems in the past. Tell your doctor if you have any of the following symptoms:

- fast or irregular heartbeat (palpitations)
- dizziness
- fainting
- chest discomfort
- shortness of breath

 **Jaypirca**[®]
pirtobrutinib 50, 100 mg tablets

A Lilly Medicine

Understanding the **possible side effects of Jaypirca**

Jaypirca can cause serious side effects, including:


- Infections
- Bleeding problems (hemorrhage)
- Decrease in blood counts (white blood cells, platelets, and red blood cells)
- Heart rhythm problems (atrial fibrillation and atrial flutter)
- Second primary cancers (new cancers, including of the skin or other organs)
- Liver problems

The most common side effects of Jaypirca include:

- Decreased white blood cell count
- Decreased red blood cell count
- Tiredness
- Decreased platelet count
- Decreased calcium

These are not all the possible side effects of Jaypirca. See accompanying Safety Summary for more information. If you experience side effects while on treatment, it's important to talk to your doctor.

9 out of 10 people with MCL kept taking Jaypirca without needing to stop treatment due to side effects



In a clinical trial:

- 9% of **people with MCL** (12 out of 128) stopped taking Jaypirca permanently due to side effects
- Side effects that caused more than 1% of patients to permanently stop taking Jaypirca included pneumonia

Jaypirca was studied in a clinical trial that included people with MCL that came back or did not respond to previous treatment, and who had already received one or more treatments for their cancer, including a BTK inhibitor medicine. The trial evaluated 120 people for their response to treatment (a response means their cancer shrank or stopped spreading) and how long the response lasted. People in the trial took Jaypirca once daily until their disease got worse or they were no longer able to tolerate treatment.

Select Important Safety Information

Warnings - Jaypirca may cause serious side effects, including:
Second primary cancers. New cancers have happened during treatment with Jaypirca, including cancers of the skin or other organs. Your doctor will check you for other cancers during treatment. Use sun protection when you are outside in sunlight.


Jaypirca[®]
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

Half of adults with **MCL** who took Jaypirca **saw their cancer shrink and stop spreading**

IN A CLINICAL TRIAL THAT MEASURED THE NUMBER OF PEOPLE WHO RESPONDED TO TREATMENT AND HOW LONG THE RESPONSE LASTED:



OF ADULTS WITH MCL (60 out of 120) saw their **cancer shrink and stop spreading** while taking Jaypirca

OF THE 50%:

13%

OF ADULTS (15 out of 120) had a complete response, which means scans and other tests showed no detectable signs of MCL

38%

OF ADULTS (45 out of 120) had a partial response, which means scans and other tests showed their MCL shrank in response to treatment

- **65% were still responding** to treatment at 6 months
- For those who saw their MCL shrink and stop spreading, their response lasted a **median* of 8.3 months**

*Median is the middle number in a range of numbers.

In a clinical trial, Jaypirca started working quickly, with some people seeing their cancer reduce in size and stop spreading as soon as 3.5 weeks after starting treatment.

In a clinical trial of Jaypirca, the median time for the cancer to shrink and stop spreading with treatment was 1.8 months, ranging from 3.5 weeks at the lowest to 4.2 months at the highest. Median is the middle number in a range of numbers ordered from lowest to highest.

Select Important Safety Information

Warnings - Jaypirca may cause serious side effects, including:

Liver Problems. Liver problems, which may be severe or life-threatening, or lead to death, can happen in people treated with Jaypirca. Your doctor will do blood tests to check your liver before and during treatment with Jaypirca. Tell your doctor or get medical help right away if you have any signs of liver problems, including stomach pain or discomfort, dark-colored urine, or yellow skin and eyes.

Your doctor may decrease your dose, temporarily stop, or permanently stop treatment with Jaypirca if you develop severe side effects.



Jaypirca[®]
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

See how you may be able to **save** on Jaypirca



Jaypirca Savings Card

Paying for treatment shouldn't be an additional concern for you and your loved ones, so we've created the Jaypirca Savings Card for eligible, commercially insured patients, which may help you manage treatment costs

Eligible, commercially insured covered patients pay as little as \$0 a month.*

*Month is defined as 30-days.

Governmental beneficiaries excluded, terms and conditions apply.

Need a Jaypirca Savings Card? Visit www.jaypirca.lilly.com/savings-support#savings for more information or call the Jaypirca Savings Card Support Line at 1-833-727-4589

By enrolling in and using the Jaypirca Savings Card Program ("Program") and using the Jaypirca Savings Card ("Card"), you attest that you meet the eligibility criteria, and you agree to comply with the terms and conditions described below:

Card Eligibility:

- (1.) You have been prescribed Jaypirca® (pirtobrutinib) for an approved use consistent with FDA-approved product labeling;
- (2.) You are enrolled in a commercial drug insurance plan and have coverage for Jaypirca;
- (3.) **You are not enrolled in any state, federal, or government funded healthcare program, including, without limitation, Medicaid, Medicare, Medicare Part D, Medicare Advantage, Medigap, DoD, VA, TRICARE®, CHAMPUS, or any state prescription drug assistance program;**
- (4.) You are a resident of the United States or Puerto Rico; and
- (5.) You are 18 years of age or older.

Card Terms and Conditions

For patients with commercial drug insurance coverage for Jaypirca: You must have commercial drug insurance that covers Jaypirca and a prescription for an approved use consistent with FDA-approved product labeling to pay as little as \$0 for a 1-month prescription fill of Jaypirca. Month is defined as 30-days. Card savings are subject to a maximum monthly savings of wholesale acquisition cost plus usual and customary pharmacy charges and separate maximum annual savings of up to \$10,600 per calendar year. Card may be used for a maximum of up to 14 prescription fills per calendar year. Except where prohibited by applicable state law, Card monthly and annual savings are reduced if Lilly identifies that you are enrolled in a plan or program, sometimes called a maximizer plan, that adjusts your cost sharing amount to be equal to or include some portion of the savings provided by the Card and attempts to prevent the savings from this Card from being applied to your out-of-pocket costs, including but not limited to copayments, coinsurances, and deductibles ("Maximizer"). If the Program identifies you are enrolled in a Maximizer, Card savings are reduced to a maximum monthly savings of up to \$25 and a separate maximum annual savings of up to \$350 per calendar year. If you have reason to believe that the Program erroneously identified enrollment in a Maximizer, please call the Jaypirca Savings Card Program at 1-833-727-4589. Subject to Lilly USA, LLC's ("Lilly") right to terminate, rescind, revoke, or amend Card eligibility criteria and/or Card terms and conditions which may occur at Lilly's sole discretion, without notice, and for any reason. Card expires and savings end on 12/31/2026.

Additional Program Terms and Conditions

If you have an insurance plan that is participating in an alternate funding program ("AFP") that requires you to apply to the Jaypirca Savings Card Program or otherwise pursue specialty drug prescription coverage through an alternate funding vendor as a condition of, requirement for,

CONTINUED ↓



Jaypirca®
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

Savings & Support

Additional Program Terms and Conditions (cont'd)

or prerequisite to coverage of Jaypirca, you are not eligible for and are prohibited from using the Jaypirca Savings Card Program. AFPs include programs where coverage, reimbursement, or patient out of pocket costs for a product in some way vary based on the availability of a manufacturer co-pay program. AFPs may modify, delay, deny, restrict, or withhold insurance benefits or coverage from patients, or exclude Lilly products from coverage contingent upon a member's use of Jaypirca Savings Card Program. You agree to inform the Jaypirca Savings Card Program if you are or become a member of such an AFP. You are responsible for any applicable taxes, fees, and any amount that exceeds the monthly or annual maximum Card savings. Monthly and annual maximum savings are set at Lilly's sole and absolute discretion and may be changed with or without notice at any time for any reason. At its sole discretion and with or without notice, Lilly may reduce, eliminate, or otherwise modify the Card savings for any reason, including but not limited to if your commercial drug insurance plan imposes additional requirements which limits or prevents you from receiving coverage for Jaypirca, only allows partial coverage for Jaypirca, removes coverage for Jaypirca and requires you to utilize the Card, does not provide a material level of financial assistance for the cost of Jaypirca, or does not apply Card payments to satisfy your co-payment, deductible, or coinsurance for Jaypirca. Card savings are not valid for: Massachusetts residents if an AB-rated generic equivalent is available; California residents if an FDA-approved therapeutic equivalent is available. You must meet the Card eligibility criteria, terms and conditions every time you use the Card. If at any time you begin receiving drug coverage under any state, federal, or government funded healthcare program, you understand that you will no longer be eligible for the Jaypirca Savings Card and agree to call the Jaypirca Savings Card Program at 1-833-727-4589 to stop participation. Card activation is required. You may not seek reimbursement from your health insurance, any third party, or any health savings, flexible spending, or other healthcare reimbursement accounts, for any amount of the savings received through the Card. By utilizing the Card, you agree that if you are required to do so under the terms of your insurance coverage for this prescription or are otherwise required to do so by law, you will notify your Insurance Carrier of your redemption of the Card. Card savings cannot be combined or utilized with any other program, discount, discount card, cash discount card, coupon, incentive, or similar offer involving Jaypirca. You agree that this Card savings is intended solely for the benefit of you, the patient, and that the Card benefits are non-transferable. It is prohibited for any person to sell, purchase, or trade; or to offer to sell, purchase, or trade, or to counterfeit the Card. **THIS CARD IS NOT INSURANCE.** Lilly has the sole right to interpret and apply Card eligibility criteria, and terms and conditions. Card eligibility, and terms and conditions may be terminated, rescinded, revoked, or amended by Lilly at any time without notice and for any reason. Lilly's sole discretion to terminate, rescind, revoke, or amend Card eligibility and/or Card terms and conditions includes the right to terminate any individual Card if Lilly determines, in its sole discretion, that a patient does not satisfy the Card's eligibility criteria or is using or has attempted to use the Card inconsistently with these terms and conditions. Eligibility criteria, and terms and conditions for the Jaypirca Savings Card Program may change from time to time; the most current version can be found at <https://www.jaypirca.lilly.com/savings-support#savings>. You may be required to obtain a new Card, including if any Card terms and conditions have been terminated, rescinded, revoked, or amended by Lilly. Card void where prohibited by law. Subject to Lilly's right to terminate, rescind, revoke or amend Card eligibility criteria and/or Card terms and conditions which may occur at Lilly's sole discretion, without notice, and for any reason. Card expires and savings end on 12/31/2026.

TRICARE® is a registered trademark of the Department of Defense (DoD), DHA.



Questions about savings?

Call the Jaypirca Savings Card Support Line at **1-833-727-4589**.

For all other questions regarding support, call Lilly Support Services™ at **1-800-LillyRx (1-800-545-5979)**.

For more information, visit **[jaypirca.lilly.com/savings-support](https://www.jaypirca.lilly.com/savings-support)**. Jaypirca support program and offerings are not a guarantee of coverage. Terms and conditions apply for all programs.



Jaypirca®
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

Safety Summary

Warnings - Jaypirca may cause serious side effects, including:

Infections. Infections can happen during treatment with Jaypirca. These infections can be serious and may lead to death. Your doctor may prescribe vaccines and certain medicines if you have an increased risk of getting infections. Tell your doctor right away if you have fever, chills, weakness, flu-like symptoms, or any other signs of infection during treatment with Jaypirca.

Bleeding problems (hemorrhage). Bleeding problems can happen during treatment with Jaypirca. They can be serious and may lead to death. Your risk of severe bleeding may increase if you are also taking a blood thinner medicine. Tell your doctor if you have any signs or symptoms of bleeding, including:

- blood in your stools or black stools (looks like tar)
- pink or brown urine
- unexpected bleeding, or bleeding that is severe or you cannot control
- vomit blood or vomit blood that looks like coffee grinds
- cough up blood or blood clots
- increased bruising
- dizziness
- weakness
- confusion
- changes in your speech
- headache that lasts a long time

Decrease in blood cell counts. Decrease in white blood cells, platelets, and red blood cells are common with Jaypirca, but can also be severe. This may increase your risk of infection, bleeding, and anemia. Your doctor should do blood tests regularly during treatment to monitor your counts.

Heart rhythm problems. Heart rhythm problems including atrial fibrillation and atrial flutter have happened in people treated with Jaypirca. Your risk for heart rhythm problems may be increased if you have high blood pressure or have had heart rhythm problems in the past. Tell your doctor if you have any of the following symptoms:

- fast or irregular heartbeat (palpitations)
- dizziness
- fainting
- chest discomfort
- shortness of breath

Second primary cancers. New cancers have happened during treatment with Jaypirca, including cancers of the skin or other organs. Your doctor will check you for other cancers during treatment. Use sun protection when you are outside in sunlight.

Liver Problems. Liver problems, which may be severe or life-threatening, or lead to death, can happen in people treated with Jaypirca. Your doctor will do blood tests to check your liver before and during treatment with Jaypirca. Tell your doctor or get medical help right away if you have any signs of liver problems, including stomach pain or discomfort, dark-colored urine, or yellow skin and eyes.

Your doctor may decrease your dose, temporarily stop, or permanently stop treatment with Jaypirca if you develop severe side effects.

CONTINUED ↓



 **Jaypirca**[®]
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

Safety Summary (cont'd)

Common side effects

- decreased white blood cell count
- decreased red blood cell count
- tiredness
- decreased platelet count
- decreased calcium

These are not all the possible side effects with Jaypirca. Call your doctor for medical advice about side effects. You may report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.

Before using

Before taking Jaypirca, tell your doctor about all your medical conditions, including if you:

- have an infection or have been advised that you are at increased risk of infection
- have had recent surgery or plan to have surgery. Your doctor may stop Jaypirca for a planned medical, surgical, or dental procedure
- have bleeding problems or are taking blood thinner medicine
- have or have had heart rhythm problems
- have high blood pressure
- have a history of other cancers, including skin cancer
- have kidney problems
- have liver problems
- are pregnant or plan to become pregnant. Jaypirca can harm your unborn baby. If you are able to become pregnant, your doctor will do a pregnancy test before starting treatment
 - Females who are able to become pregnant should use effective birth control (contraception) during treatment and for at least 1 week after the last dose of Jaypirca. Tell your doctor right away if you become pregnant or think you are pregnant during treatment
- are breastfeeding or plan to breastfeed. It is not known if Jaypirca passes into breast milk. Do not breastfeed during treatment and for at least 1 week after your final dose

Also tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking Jaypirca with certain other medicines may affect how Jaypirca or the other medicines work and can cause side effects.

CONTINUED ↓



Jaypirca[®]
pirtobrutinib 50, 100mg
tablets

Safety Summary (cont'd)

How to take

- Take Jaypirca exactly as your doctor tells you
- Do not change your dose or stop taking Jaypirca unless your healthcare provider tells you to do so
- Take Jaypirca 1 time each day at about the same time each day
- Take Jaypirca with or without food
- Swallow Jaypirca whole with water. Do not cut, crush, or chew the tablets
- If you miss a dose of Jaypirca, take it as soon as you remember on the same day. If you miss taking a dose by more than 12 hours from the time you usually take Jaypirca, skip the missed dose. Take the next dose on the next day at your usual time

Learn more

Jaypirca is a prescription medicine. For more information, call 1-800-545-5979 or go to www.jaypirca.lilly.com. This summary provides basic information about Jaypirca but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other healthcare provider about Jaypirca and how to take it. Your doctor is the best person to help you decide if Jaypirca is right for you.

PT CON BS MCL DEC2025



Not an actual patient.

ASK YOUR DOCTOR ABOUT JAYPIRCA

Please see [Safety Summary](#) for Jaypirca on pages 8-10.


Jaypirca[®]
pirtobrutinib 50, 100mg
tablets

A Lilly Medicine

Lilly

A MEDICINE COMPANY

Jaypirca[®] is a registered trademark and Lilly Support Services[™] is a trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates. All other trademarks are the property of their respective owners. PP-PT-US-1426 12/2025 © Lilly USA, LLC 2025. All rights reserved.